

# **Post-Operative Care Instructions for Thyroid and Parathyroid Surgery**

#### **Wound Care**

- A **special skin glue** was applied over your incision to protect it from dirt and water.
  - o The glue may gradually peel off—do not pull on it, but you may trim loose edges.
  - Any remaining glue will be removed at your follow-up visit.
- **Showering is allowed**, but avoid scrubbing the incision or submerging it in water (baths, pools, or the ocean).
- All **sutures** are absorbable and will dissolve naturally—no removal needed.
- **Protect the incision from sunlight** until fully healed (typically **3-4 months**, but in some cases, up to a year).
  - Cover the incision or apply SPF 30+ sunscreen to prevent darkening (hyperpigmentation).

### **Activity**

- It is normal to feel **tired** and have a **decreased appetite** on the day of surgery—rest is essential.
- Some patients feel better than expected, but it is important to take things easy for the first week.
  - Avoid straining, physical exertion, or lifting over 5 kg, as these activities increase blood pressure and risk of bleeding.
- No swimming or activities that fully submerge the incision until cleared by your surgeon.

## **Pain Management**

- Neck soreness and bruising are common after surgery.
  - As the incision heals, it may itch or burn.
  - For discomfort, take paracetamol and ibuprofen/celecoxib either together or staggered—this is more effective than narcotics.
- Throat soreness, dryness, or scratchiness is common due to the breathing tube used during anesthesia.
  - These symptoms improve within a week.
  - You may use throat sprays or lozenges for relief.

#### Medications

- Some patients require thyroid hormone replacement, calcium, or vitamin D after surgery.
  - o If prescribed, take them as directed—they are essential for your health.
  - Not all patients need these medications—your surgeon will determine what is necessary and prescribe these for you.

## **Voice Changes**

- Your voice may sound **different** after surgery due to irritation of the muscles around the voice box.
  - o This usually improves over a few weeks.
  - Sometimes, voice pitch may permanently change slightly due to muscle adjustments during surgery.
  - If lingering voice concerns persist, your surgeon may use a scope to visualize your vocal cords.

### When to Call Your Surgeon

Contact us **immediately** if you experience:

- Fever over 38.5°C.
- Sudden swelling around the incision.
- Sudden increase in pain at the incision site.
- Pus drainage from the incision.
- **Numbness or tingling** around the **mouth or fingers**. If this occurs take an extra dose of calcium supplements and contact your surgeon.
- · Any symptoms that seem unusual or concerning

## Follow-Up Care

- Your first **follow-up visit** will typically be within **2 weeks** after surgery.
  - The skin glue will be removed, and your incision will be evaluated.
- Pathology results take 7-10 days to return.

### **Contact Information**

- Non-Urgent Questions: Call the office at 03 548 4994.
- **Urgent Concerns:** Contact Dr. Slough directly. Leave a message or text if there is no immediate response, especially after hours.